

GRAVITATIONAL SETTLING OF ²²Ne IN LIQUID WHITE DWARF INTERIORS

LARS BILDSTEN

Institute for Theoretical Physics and Department of Physics
 Kohn Hall, University of California, Santa Barbara, CA 93106
 bildsten@itp.ucsb.edu

AND

DAVID M. HALL

Department of Physics, Broida Hall, University of California, Santa Barbara, CA 93106
 halldm@physics.ucsb.edu

To Appear in Astrophysical Journal Letters

Draft version February 1, 2008

ABSTRACT

The nuclear reactions that occur in the stellar progenitors of white dwarfs (WDs) lead to an internal composition of ¹²C, ¹⁶O, and a “contaminant” nucleus, ²²Ne. The ²²Ne is produced by helium captures on ¹⁴N left from hydrogen burning via the CNO cycle. By virtue of its two excess neutrons (relative to the predominant $A = 2Z$ nuclei), a downward force of $\approx 2m_p g$ is exerted on ²²Ne in the WD interior. This biases its diffusive equilibrium, forcing ²²Ne to settle towards the center of the WD. We discuss the physics of the gravitational settling when the WD is in the liquid state and the luminosity generated by it. This modifies the cooling of WD’s with masses in excess of $M_\odot$. The current uncertainties in the microphysics even allow for solutions where a $1.2M_\odot$ WD remains mostly liquid for a few Gyrs due to the internal heating from ²²Ne sedimentation. This highlights the need for an accurate calculation of the inter-diffusion coefficient, especially in the quantum liquid regime relevant for high mass WD’s. There is also time in old, liquid WD’s (such as those found in cataclysmic variables and possibly in accreting Type Ia progenitors) for partial settling.

Subject headings: diffusion – novae, cataclysmic variables – stars: abundances, interiors – supernovae: general – white dwarfs

1. INTRODUCTION

The slowest step in the CNO cycle is the proton capture on ¹⁴N. This results in all CNO catalysts piling up at ¹⁴N when H burning is completed during the main sequence. During Helium burning, the reactions ¹⁴N(α, γ)¹⁸F(β^+)¹⁸O(α, γ)²²Ne ensue and convert the ¹⁴N into ²²Ne, resulting in a mass fraction of ²²Ne, $X_{22} \approx Z_{CNO} \approx 0.02$ for Population I. The possibility that this nucleus plays an important role in the energetics of cooling WD’s was first noted by Isern et al. (1991). They discussed the possibility of gravitational energy release if ²²Ne phase separates at crystallization. This was followed by more detailed studies (Xu & Van Horn 1992; Ogata et al. 1993; Segretain et al. 1994; Segretain 1996) which differed in their conclusions about the final state once crystallization was complete; ranging from all the ²²Ne in the stellar center to most of it having an unchanged profile.

What about ²²Ne sedimentation in the liquid state? Ogata et al. (1993) showed that demixing in the liquid state never occurs for the C/O/²²Ne liquid (namely it will remain in the fully mixed state when the temperature is safely above melting). This allows us to focus on the gravitational settling of individual ²²Ne nuclei in the liquid parts of the white dwarf (Bravo et al. 1992). This happens prior to the onset of crystallization in a cooling WD and during the lifetime of a WD that remains in the liquid state due to heating from accretion.

Degenerate electrons dominate the pressure in the WD interior, so that the upward pointing electric field at mass coordinate $m(r)$ is $eE \approx 2m_p g$, where $g = Gm(r)/r^2$ and m_p is the proton mass. The downward force on a ²²Ne nucleus is $F = 22m_p g - 10eE = 2m_p g$, giving a potential energy drop $U(r) = \int_0^r 2m_p g dr$ between radius r and the center. Across

the whole star, this amounts to 285 keV for $M = 0.6M_\odot$ and 1600 keV for $M = 1.2M_\odot$. It is the excess neutrons of the ²²Ne nucleus (relative to $A = 2Z$) that make it special (the same is true of the less abundant ⁵⁶Fe). Differentiating forces on the carbon and oxygen nuclei only arise from the slight differences in masses from Am_p due to the nuclear binding, which is orders of magnitude smaller than the force on ²²Ne.

The potential energy (the energy that would be released if all the ²²Ne sinks to the center) of a uniformly mixed model ($X_{22} = 0.02$ is constant) is 1.4×10^{47} , 3.2×10^{47} , 6.8×10^{47} and 1.5×10^{48} erg for masses of 0.6, 0.8, 1.0, & $1.2M_\odot$. For 0.6 and $1.0M_\odot$, this corresponds to 5.8×10^{15} & 1.7×10^{16} erg gr⁻¹ of possible energy release from ²²Ne settling. At a temperature of 10^7 K, the specific heat per oxygen ion is 7.7×10^{13} erg gr⁻¹, so that the energy stored in ²²Ne is comparable to the thermal energy content of a cooling WD at late times.

In §2, we estimate the inter-diffusion coefficient for a trace ²²Ne nucleus in the liquid C/O interior. The settling timescale for ²²Ne is ~ 10 Gyr, making it critical for a more accurate calculation of the inter-diffusion coefficient. The settling time decreases for more massive WDs. In §3, we speculate on the modifications the ²²Ne settling is likely to have on accreting WD’s and maybe Type Ia supernovae. We calculate the luminosity from the gravitational energy release of sinking ²²Ne in §4. *This affects the cooling of WD’s prior to crystallization and can possibly heat them to a level hot enough to maintain them in the liquid state for a Hubble time.*

2. DIFFUSION AND SETTLING OF THE ²²Ne

We presume that the WD interior consists of a single element (either ¹²C or ¹⁶O) of mass Am_p and charge Ze , and that ²²Ne is

a trace component at density $n_{\text{Ne}} = X_{22}\rho/22m_p$. Our WD is constructed purely from the degenerate electron equation of state. For the properties we need for the ^{22}Ne diffusion problem, it is a fine approximation to neglect the ionic contribution to the pressure.

Relative diffusion of charged species in a correlated liquid is substantially different than the diffusion problem in the WD atmosphere. In those less dense regions, traditional Coulomb cross sections and ideal gas equations of state dominate (e.g. Fontaine & Michaud 1979; Vauclair & Vauclair 1982). As discussed by Paquette et al. (1986), there is substantial uncertainty in the diffusion coefficients in the regions of the WD where the ion Coulomb coupling becomes strong. For a classical one component plasma (OCP) with ion separation, a , defined by $a^3 = 3/4\pi n_i$, where $n_i = \rho/Am_p$, the importance of Coulomb physics for the ions is measured by

$$\Gamma \equiv \frac{(Ze)^2}{akT} = 57.7\rho_6^{1/3} \left(\frac{10^7 \text{ K}}{T}\right) \left(\frac{Z}{8}\right)^2 \left(\frac{16}{A}\right)^{1/3}, \quad (1)$$

where $\rho_6 = \rho/10^6 \text{ gr cm}^{-3}$. Crystallization occurs when Γ exceeds 173 (see Farouki & Hamaguchi 1993 and references therein). We only consider settling of ^{22}Ne in the liquid state, or $1 < \Gamma < 173$, the presumption being that the dramatic increase in viscosity expected in the solid state will prohibit further gravitational sedimentation.

The inter-diffusion coefficient, D , of ^{22}Ne in a C/O mixture with $\Gamma \sim 1 - 100$ has not been specifically calculated and so we construct here our best estimate. One way to estimate D is to use the Stokes-Einstein relation for a particle of radius a_p (in this case ^{22}Ne) undergoing Brownian motion in a fluid of viscosity η , which gives $D = kT/4\pi a_p \eta$ when the fluid is allowed to “slip” at the particle/fluid interface. This estimate works well in other liquids at small scales, where it is accurate at atomic dimensions (Hansen & McDonald 1986). An alternate route is to use the self-diffusion coefficient in the OCP

$$D \approx 3\omega_p a^2 \Gamma^{-4/3}, \quad (2)$$

calculated by Hansen, McDonald and Pollock (1975), where $\omega_p^2 = 4\pi n_i (Ze)^2 / Am_p$ is the ion plasma frequency. How well does this agree with the Stokes-Einstein estimate? Using the numerically derived viscosity of the OCP, $\eta \approx 0.1\rho\omega_p a^2 (\Gamma/10)^{0.3}$ (Donko & Nyiri 2000 and references therein) and setting $a = a_p$, the derived $D = kT/4\pi a \eta$ agrees with equation (2) to about 20% for all $\Gamma > 10$.

We also compared equation (2) to Fontaine’s (1987) rough fitting formula for the interdiffusion coefficients and found agreement to $\approx 30\%$. So, for now, we will use equation (2) for our estimate of D , keeping in mind that there has yet to be an appropriate microphysical calculation. We will show later that there is astrophysical motivation for a more accurate calculation of the inter-diffusion coefficients, especially in the quantum liquid regime that is relevant for the massive WD’s. For these WDs, the parameter $\hbar\omega_p/kT$ exceeds unity long before crystallization sets in (Chabrier, Ashcroft & DeWitt 1992; Chabrier 1993) and might well modify the diffusion coefficient by order unity.

¹Our present understanding of the microphysics does not predict the development of a pure ^{22}Ne core. However, if this were to occur, it is unlikely to modify the ignition conditions, as the electron energy threshold for the reaction $^{22}\text{Ne} + e^- \rightarrow ^{22}\text{F} + \nu_e$ is 11.36 MeV, which requires a density in excess of $2.1 \times 10^{10} \text{ g cm}^{-3}$. Carbon ignites at densities about an order of magnitude less (Nomoto, Thielemann & Yokoi 1984), so that the initial trigger of carbon burning is unchanged. However, a pure ^{22}Ne core with a mass of $0.028M_\odot$ would have a radius of $R_{\text{Ne}} \approx 190 \text{ km}$ at a density of $2 \times 10^9 \text{ g cm}^{-3}$, forcing the C/O mixture to ignite in the shell covering the ^{22}Ne core.

Since $kT \ll U(r)$ over most of the stellar radius (equivalent to saying that the ^{22}Ne scale height is much less than the WD radius), most of the drift of the ^{22}Ne is dominated by “falling” at the local speed set by the diffusion coefficient $V \approx 2m_p gD/kT$, which with equation (2) gives

$$V = \frac{18m_p g}{Ze\Gamma^{1/3}(4\pi\rho)^{1/2}}. \quad (3)$$

In mass coordinates, $m(r)$, the time it takes to fall from m_1 to m_2 is

$$\Delta t = \int_{m_1}^{m_2} \frac{Ze\Gamma^{1/3}}{18m_p G(4\pi\rho)^{1/2}} \frac{dm(r)}{m(r)}. \quad (4)$$

For a constant density star, this becomes $\Delta t \approx t_s \int d \ln m(r)$, where

$$t_s \equiv \frac{\Gamma^{1/3}Z}{18} \left(\frac{e^2}{Gm_p^2}\right)^{1/2} \left(\frac{1}{4\pi G\rho}\right)^{1/2} = 13 \text{ Gyr} \frac{Z\Gamma^{1/3}}{6\rho_6^{1/2}}. \quad (5)$$

Using the central densities as an estimate, a pure oxygen $0.6M_\odot$ WD at $T = 10^8 \text{ K}$ (this temperature is appropriate for rapidly accreting WD’s) has $t_s \approx 19.6 \text{ Gyr}$, so that complete sedimentation is not likely for the most common $0.6M_\odot$ WD’s. However, the strong density dependence means that this settling time decreases for massive WD’s. For example, a pure oxygen $1.2(1.3)M_\odot$ WD at 10^8 K has $t_s \approx 4.8(3) \text{ Gyr}$, in rough agreement with the previous simple estimates of Bravo et al. (1992).

Figure 1 shows the time it takes for a ^{22}Ne nucleus that starts at the surface ($m(r) = M$) to fall to the location, $m(r)$. This is an integration of equation (4) with a model WD. The solid (dashed) lines are for pure oxygen (carbon) and bracket the C/O WD’s. We show two masses, 0.6 and $1.2 M_\odot$, making it clear that ^{22}Ne settles faster in a massive WD.

3. ACCRETING WHITE DWARFS AND TYPE IA PROGENITORS

Unlike cooling WDs, the interiors of massive, accreting WD’s are maintained in the liquid state by the heating from accretion. For slowly accreting ($\dot{M} \sim 10^{-10} M_\odot \text{ yr}^{-1}$) WD’s in cataclysmic variables below the period gap, the core temperatures are in the range $T \approx (1-3) \times 10^7 \text{ K}$ (Nomoto 1982), so that they remain liquid while accreting for a few Gyrs. A $1M_\odot$ pure oxygen WD in such a setting will undergo neon settling in the outer layers as $t_s \approx 12.6 \text{ Gyr}$ at $T = 3 \times 10^7 \text{ K}$.

Though the debate rages regarding appropriate Type Ia progenitor models, most favor rapidly accreting WDs that ignite carbon once the mass is near the Chandrasekhar value. The accretion rates considered are too rapid to allow ^{22}Ne settling during this phase of accretion. However, prior evolution at lower accretion rates or an extended period as a cooling WD will have allowed for some Neon settling. Though unclear just how important this will be, the effects of ^{22}Ne settling for the Type Ia ignition of a near-Chandrasekhar mass WD are important to note. The most obvious one is that ^{22}Ne settling will develop a gradient in the electron mean molecular weight. This would modify the convective criterion in the ignited core and possibly affect the evolution from the thermal runaway to a propagating flame front (Garcia-Senz & Woosley 1995).¹

The ^{22}Ne abundance in a Type Ia progenitor is also critical to the overproduction of the neutron rich isotopes ^{54}Fe and ^{58}Ni (Thielemann, Nomoto & Yokoi 1986). The neutron excess in the inner parts of an exploding WD is fixed by electron captures. Modifying the production of these elements at that location would require a substantially enhanced ^{22}Ne abundance there (e.g. a pure ^{22}Ne core would set the initial $Y_e \approx 0.46$). Outside of $(0.3-0.4)M_\odot$, the production of ^{54}Fe and ^{58}Ni is fixed by the local ^{22}Ne abundance (Iwamoto et al. 1999). Hence, the production of these elements can be reduced if the ^{22}Ne sinks away from the outer regions (Bravo et al. 1992).² Our calculation suggests there is time for ^{22}Ne settling in the outer liquid layers of a massive WD if accreting at a low rate. The final answer to this puzzle awaits a time-dependent evolution on a cosmological timescale of the diffusion equation with a realistic starting abundance profile and accretion history.

There is also a ^{22}Ne settling calculation relevant to classical novae. Many lines of evidence (ejected masses and overly abundant heavy elements) point to the ejected material in classical novae containing matter dredged up from the underlying C/O WD (see Gehrz et al. 1998 and Starrfield 1999 for overviews). Livio & Truran (1994) noted that the Neon abundances measured in many of these ejecta are consistent with the $X_{22} \approx 0.02$ expected in a C/O WD. Hence, there is no need to invoke the presence of an O/Ne/Mg WD for those events. However, this will remain true only if the ^{22}Ne has not gravitationally settled out of the outer C/O layers of the WD that are about to be dredged up.

We can assess this by finding how far the ^{22}Ne falls in the surface C/O layers, where g is constant. This is a simple case, where we integrate equation (4) for an isothermal layer. Using pressure as a coordinate, we find that the time it takes for ^{22}Ne to fall to a pressure P starting from the surface is

$$\Delta t \approx 0.8 \text{ Gyr } \Gamma_p^{1/3} \left(\frac{Z}{6} \right) \left(\frac{10^8 \text{ cm s}^{-2}}{g} \right)^2 \left(\frac{P}{10^{20} \text{ erg cm}^{-3}} \right)^{0.7} \quad (6)$$

where Γ_p is the value of Γ at pressure P and exceeds one at the time of ignition.

Since the accreted material appears to dredge up a WD layer of mass comparable to the accreted layer, we just need to compare the ^{22}Ne settling time to the time it takes to accumulate the fuel, $t_a = 4\pi GMP/g^2\dot{M}$. The ratio of these timescales gives

$$\frac{\Delta t}{t_a} \approx 150 \Gamma_p^{1/3} \left(\frac{M_\odot}{M} \right) \left(\frac{\dot{M}}{10^{-9} M_\odot \text{ yr}^{-1}} \right) \left(\frac{2 \times 10^{19} \text{ erg cm}^{-3}}{P} \right)^{0.3}, \quad (7)$$

where we have written the pressure in units of the typical ignition pressure (Livio 1994). Thus, for $\dot{M} > 10^{-11} M_\odot \text{ yr}^{-1}$ (which applies to most classical novae, Livio 1994), the underlying C/O material should have ^{22}Ne present at $X_{22} \approx 0.02$, *allowing for neon enrichment on a C/O WD in those classical novae that are constantly undergoing dredge-up.*

4. LUMINOSITIES FROM FALLING ^{22}Ne IN COOLING WHITE DWARFS: SEDIMENTARS

The rate of energy release from the falling ^{22}Ne is given by the integral of the power, FV ,

$$L_g \equiv \int FV n_{\text{Ne}} 4\pi r^2 dr. \quad (8)$$

Since the ^{22}Ne settling time is longer than the time it takes to cool to the onset of crystallization in the core (about $2(0.3)$ Gyr for a $0.6(1.2)M_\odot$ WD; Benvenuto & Althaus 1999), we presume that the abundances are nearly unchanged from the initial state. This integral is evaluated with $X_{22} = 0.02$ and the core temperature, T_c , being constant. For a pure oxygen WD at $T_c = 10^8$ K, this gives $L_g = 1.1 \times 10^{29}, 3.9 \times 10^{29}, 1.2 \times 10^{30} \& 4.6 \times 10^{30} \text{ erg s}^{-1}$ for $M = 0.6, 0.8, 1.0, \& 1.2 M_\odot$.

For massive WD's, L_g becomes comparable to the cooling luminosity, L , for the same T_c . This raises the important question of how much the settling luminosity can affect the WD cooling curves and thus the inferred age of the galactic disk from the faint end of the WD luminosity function (see Leggett, Ruiz and Bergeron 1998 for a recent overview).

We begin with the common $0.6M_\odot$ WDs. The hatched region in the top panel of Figure 2 shows L_g for $X_{22} = 0.02$ as a function of T_c . The upper (lower) bound of the hatched region is for a pure carbon (oxygen) WD and denotes the range of possibilities for arbitrary C/O mixtures. The sharp downturn at $T_c \approx (3-4) \times 10^6 \text{ K}$ is due to core crystallization. We only allow for sedimentation in the liquid parts of the WD, so that, as T_c decreases and the solid core increases, L_g decreases. If we allowed for continuing sedimentation throughout the solidifying WD, the hatched region would just continue to the left with the same slope. The dashed line is the $L - T_c$ relation for a cooling $0.6M_\odot$ WD (see caption). For $0.6M_\odot$, L_g is always at least a factor of 10 lower than the exiting luminosity, L , at that T_c .

However, this competition gets closer as the WD mass increases. The bottom panel of Figure 2 shows the same curves for $1.2M_\odot$. In this case, L_g is a factor of 5 or so less than L , which means that the WD cooling trajectory would be affected considerably. It is a coincidence that L_g drops off before becoming dominant and then remains nearly parallel to the $L - T_c$ relation. This would not occur in a pure helium WD, as crystallization never sets in. However, in that case, there would be no ^{22}Ne present. It would just be the settling of the other, less abundant, neutron rich metals, such as ^{56}Fe . Though it appears unlikely that the settling of iron will modify the cooling of a low-mass helium ($M < 0.4M_\odot$) WD, substantial settling will occur in 10 Gyr.

The point we wish to make here is that the uncertain microphysics of the diffusion coefficient clearly allows for an alternative view of cooling for massive WD's. Namely, it is possible that the settling luminosity can heat a white dwarf well enough to keep it in the liquid state for a very long time. For example, increasing D by a factor of 6 for the $1.2M_\odot$ WD would shift the L_g curves up enough so that the WD could be powered by sedimentation alone: hence, our dubbing of these solutions as "Sedimentars". Or, if we allow for continued sedimentation once the core is supposedly solid (say, it is supercooled), the lines would intersect at $L \approx 2 \times 10^{30} \text{ erg s}^{-1}$ for a $1.2M_\odot$ WD, which would then take a time in excess of 10 Gyr to radiate all of the sedimentation energy. This sedimentar would live for many Gyr's at an effective temperature of 11,350 K. For a pure oxygen $1.36M_\odot$ WD this crossover would occur at $L \approx 10^{31} \text{ erg s}^{-1}$ or $T_{\text{eff}} \approx 21,000 \text{ K}$, still too cold to help explain the possible excess of hot, massive WD's reported by Vennes (1999).

²Höflich et al. (1998, 2000) studied the differences in Type Ia spectral evolution that are possible if the ^{54}Fe was absent in the outer layers. These can be large in the UV, where the iron is a major opacity source. Lentz et al. (2000) varied the ^{54}Fe abundance in the partially burned outer layer of model W7 to see if the observations can accommodate a varying value. Though they also found spectral differences in the UV, nothing large enough to allow them to constrain the ^{22}Ne abundance.

5. CONCLUSIONS

Our initial calculations suggest a number of important problems to address in the future. The coincidence of the settling time with the Hubble time forces us to reconsider the physics of the inter-diffusion coefficients, especially in the quantum liquid realm. This is critically important for the cooling of massive WD's where there is some chance for the WD to shine for Gyrs on the power of ^{22}Ne sedimentation. These gravitationally powered "sedimentars" would be a new astrophysical object that behave differently than the conventional cooling WD.

The time-dependent evolution of the ^{22}Ne abundance in different astrophysical settings is the subject of our future work. In that paper, we evolve both cooling and accreting WD's with active ^{22}Ne settling. This will allow for an accurate calculation of the evolution of the ^{22}Ne abundance from starting conditions that reflect the prior stellar evolution. For cooling WD's, we can follow the ^{22}Ne throughout the star as it crystallizes, and assess the possibility of altering the nature of the phase separation at freezing via enhanced ^{22}Ne abundances.

The downward diffusion speed is very slow and it is natural to question whether such slow speeds are actually relevant. We believe so, as there is no expectation for active convection to undo it. Comparably weak diffusive settling also appears in the Sun, where inclusion of the slow gravitational settling of helium dramatically improves the agreement of the solar interior model with that measured by seismological inversions (Christensen-Dalsgaard et al. 1996; Basu, Pinsonneault & Bahcall 2000).

For conversations about this problem, we thank N. Ashcroft, D. Chernoff, S. Kulkarni, J. Liebert, E. Salpeter, and D. Stevenson. For discussions about current white-dwarf cooling models, we thank B. Hansen and D. Townsley. O. Straniero provided information on the ^{22}Ne distribution in the WD's and the anonymous referee's comments greatly improved our discussion. This research was partially supported by NASA via grant NAG 5-8658 and by the National Science Foundation under Grants PHY94-07194, PHY99-07949 and AY97-31632. L. B. is a Cottrell Scholar of the Research Corporation.

REFERENCES

- Basu, S., Pinsonneault, M. H. & Bahcall, J. N. 2000, *ApJ*, 529, 1084
 Benvenuto, O. G. & Althaus, L. G. 1999, *MNRAS*, 303, 30
 Bravo, E., Isern, J., Canal, R. & Labay, J., 1992, *A & A*, 257, 534
 Chabrier, G. 1993, *ApJ*, 414, 695
 Chabrier, G., Brassard, P., Fontaine, G., & Saumon, D. 2000, *ApJ*, 543, 216
 Chabrier, G., Ashcroft, N. W., & DeWitt, H. E. 1992, *Nature*, 360, 48
 Christensen-Dalsgaard, J. et al. 1996, *Science*, 272, 1286
 Donko, Z. & Nyiri, B. 2000, *Phys. of Plasmas*, 7, 45
 Farouki, R. T. & Hamaguchi, S. 1993, *Phys. Rev. E.*, 47, 4330
 Fontaine, G. 1987, in "Strongly Coupled Plasma Physics", ed. F. J. Rogers and H. E. Dewitt (Plenum Press: New York), p. 161
 Fontaine, G. & Michaud, G. 1979, *ApJ*, 231, 826
 Garcia-Senz, D. & Woosley, S. E. 1995, *ApJ*, 454, 895
 Gehrz, R. D., Truran, J. W., Williams, R. E. & Starrfield, S. 1998, *PASP*, 110, 3
 Hansen, B.M.S. 1999, *ApJ*, 520, 680
 Hansen, J.-P., McDonald, I. R., & Pollock, E. L. 1975, *Phys. Rev.*, A11, 1025
 Hansen, J.-P. & McDonald, I. R. 1986, *Theory of Simple Liquids* (Academic Press: London)
 Höflich, P., Wheeler, J. C. & Thielemann, F.-K. 1998, *ApJ*, 495, 617
 Höflich, P., Nomoto, K., Umeda, H. & Wheeler, J. C. 2000, *ApJ*, 528, 617
 Isern, J., Mochkovitch, R., Garcia-Berro, E., & Hernanz, M. 1991, *A & A*, 241, L29
 Iwamoto, K., Brachwitz, F., Nomoto, K., Kishimoto, N., Umeda, H., Hix, W. R., & Thielemann, F. K. 1999, *ApJS*, 125, 439
 Leggett, S. K., Ruiz, M. T. & Bergeron, P. 1998, *ApJ*, 497, 294
 Lentz, E. J., Baron, E., Branch, D., Hauschildt, P. H. & Nugent, P. E. 2000, *ApJ*, 530, 966
 Livio, M. 1994 in "Interacting Binaries" ed. S. N. Shore, M. Livio and E. P. J. van den Heuvel (Springer-Verlag: Berlin), p. 135
 Livio, M. & Truran, J. W. 1994, *ApJ*, 425, 797
 Nomoto, K. 1982, *ApJ*, 253, 798
 Nomoto, K., Thielemann, F.K., & Yokoi, K. 1984, *ApJ*, 286, 644
 Ogata, S., Iyetomi, H., Ichimaru, S. & Van Horn, H., 1993, *Phys. Rev.*, E48, 1344
 Paquette, C., Pelletier, C., Fontaine, G. & Michaud, G. 1986, *Ap. J. Suppl.*, 61, 177
 Segretain, L., Chabrier, G., Hernanz, M., Garcia-Berro, E., Isern, J., & Mochkovitch, R. 1994, *ApJ*, 434, 641.
 Segretain, L. 1996, *A & A*, 310, 485
 Starrfield, S. 1999, *Phys. Rep.*, 311, 371
 Thielemann, F.-K., Nomoto, K. & Yokoi, K. 1986, *A&A*, 158, 17
 Vauclair, S. & Vauclair, G. 1982, *ARAA*, 20, 37
 Vennes, S. 1999, *ApJ*, 525, 995
 Xu, Z. W. & Van Horn, H. M. 1992, *ApJ*, 387, 662




FIG. 1.— The gravitational settling time for a ^{22}Ne nucleus in a pure C (dashed) or pure O (solid) WD at a temperature of 10^8 K appropriate for a Type Ia progenitor. The lines show how long it takes a ^{22}Ne nucleus to fall to a given mass coordinate if it begins at the stellar surface. The set that begins at 1.2 (0.6) $M_{\odot}$ is for a WD of that mass.




FIG. 2.— The luminosity from ^{22}Ne settling in a $0.6M_{\odot}$ (top panel) and $1.2M_{\odot}$ (bottom panel) WD with $X_{22} = 0.02$. The hatched region shows the luminosity from ^{22}Ne settling as a function of WD core temperature. The upper (lower) bound is for a pure carbon (oxygen) WD and should bracket the range of possible C/O mixtures. The dashed lines are the $T_c - L$ relations for cooling DA WD's from Benvenuto & Althaus (1999). Both have helium layers of $M_{\text{He}} = 10^{-2}M_{\odot}$, while the $0.6(1.2)M_{\odot}$ WD has a hydrogen layer mass of $10^{-4}(10^{-6})M_{\odot}$. The $0.6M_{\odot}$ model is close to that of Hansen (1999) and Chabrier et al. (2000).